Royal College of Surgeons in Ireland

repository@rcsi.com

Transcriptional response of polycomb group genes to status epilepticus in mice is modified by prior exposure to epileptic preconditioning.

AUTHOR(S)

James P. Reynolds, Suzanne FC Miller-Delaney, Eva M. Jimenez-Mateos, Takanori Sano, Ross C. McKiernan, Roger P. Simon, David C. Henshall

CITATION

Reynolds, James P.; Miller-Delaney, Suzanne FC; Jimenez-Mateos, Eva M.; Sano, Takanori; McKiernan, Ross C.; Simon, Roger P.; et al. (2015): Transcriptional response of polycomb group genes to status epilepticus in mice is modified by prior exposure to epileptic preconditioning.. figshare. Journal contribution. https://hdl.handle.net/10779/rcsi.10792226.v1

HANDLE

10779/rcsi.10792226.v1

LICENCE

CC BY-NC-ND 4.0

This work is made available under the above open licence by RCSI and has been printed from https://repository.rcsi.com. For more information please contact repository@rcsi.com

URL

https://repository.rcsi.com/articles/Transcriptional_response_of_polycomb_group_genes_to_status_epilepticus_in_mice_is_modified_by_prior_exposure_to_epileptic_preconditioning_/10792226/1

Supplementary Figure 1 | Sham surgery does not alter basal Polycomb Group transcription in the hippocampus. Hippocampal homogenates were analysed by qRT-PCR and the ΔCT for various core PcG transcripts against an endogenous control, β-actin, was plotted. Measurements were from sham-surgery mice receiving intraamygdala injection of PBS (n = 4 for each timepoint) with analysis performed on microdissected hippocampal fractions enriched for **(a)** CA3, **(b)** CA1 and **(c)** DG. CA, cornu ammonis; CON, control; DG, dentate gyrus.; n.d., no data.