



repository@rcsi.com

National Audit of Hospital Mortality Annual Report 2018

AUTHOR(S)

Bridget Egan, Simon Jones, Declan McKeown, Alan Egan, Deirdre Burke, Aisling Connolly, Fionnola Kelly, The National Audit of Hospital Mortality Governance Committee, The National Office of Clinical Audit (NOCA)

CITATION

Egan, Bridget; Jones, Simon; McKeown, Declan; Egan, Alan; Burke, Deirdre; Connolly, Aisling; et al. (2021): National Audit of Hospital Mortality Annual Report 2018. Royal College of Surgeons in Ireland. Report. https://doi.org/10.25419/rcsi.16964788.v1

DOI

10.25419/rcsi.16964788.v1

LICENCE

CC BY-NC-SA 4.0

This work is made available under the above open licence by RCSI and has been printed from https://repository.rcsi.com. For more information please contact repository@rcsi.com

URL

https://repository.rcsi.com/articles/report/National_Audit_of_Hospital_Mortality_Annual_Report_2018/1696478 8/1





NATIONAL AUDIT OF HOSPITAL MORTALITY SUPPORTING APPENDIX 2018





National Audit of Hospital Mortality SUPPORTING APPENDIX 2018

This document is supplemental to the National Audit of Hospital Mortality Annual Report 2018.

For further information please refer to the main report available at https://www.noca.ie/publications/publications-listing/PO/category/3

CONTENTS

METHODOLOGY FOR MEASURING IN-HOSPITAL MORTALITY	6
FRAMEWORK FOR THE NAHM ANNUAL REPORT	9
NAHM GOVERNANCE STRUCTURE	10
INDICATORS FOR KEY DIAGNOSIS AND TABULAR INFORMATION TO SUPPORT FUNNEL PLOTS IN THE MAIN REPORT	12
ACUTE MYOCARDIAL INFARCTION	12
HEART FAILURE	14
ISCHAEMIC STROKE	16
HAEMORRHAGIC STROKE	18
COPD	20
PNEUMONIA	22

TABLES

TABLE 1	Criteria for selection of key diagnoses	9
TABLE 2	Attendance at NAHM governance committee meetings, 2018	11
TABLE 3	AMI indicator	12
TABLE 4	Tabular presentation for AMI in-hospital mortality, 2018	13
TABLE 5	Heart failure indicator	14
TABLE 6	Tabular presentation for heart failure in-hospital mortality, 2018	15
TABLE 7	Ischaemic stroke indicator	16
TABLE 8	Tabular presentation for ischaemic stroke in-hospital mortality, 2018	17
TABLE 9	Haemorrhagic stroke indicator	18
TABLE 10	Tabular presentation for haemorrhagic stroke in-hospital mortality, 2016-2018	19
TABLE 11	COPD indicator	20
TABLE 12	Tabular presentation for COPD in-hospital mortality, 2018	21
TABLE 13	Pneumonia indicator	22
TABLE 14	Tabular presentation for pneumonia in-hospital mortality, 2018	23

FIGURES

FIGURE 1	Concept of direct standardisation	7
FIGURE 2	NAHM governance structure	10

GLOSSARY

ACRONYM	FULL TERM
АМІ	acute myocardial infarction
HSE	Health Service Executive
ICD-10-AM	International Statistical Classification of Diseases and Related Health Problems, 10th Revision, Australian Modification
NAHM	National Audit of Hospital Mortality
NOCA	National Office of Clinical Audit
NQAIS	National Quality Assurance Improvement System. A suite of audit and performance-monitoring tools developed by the Health Intelligence Unit, Strategic Planning and Transformation, HSE.
NQAIS NAHM	National Quality Assurance Improvement System, National Audit of Hospital Mortality web-based tool
OECD	Organisation for Economic Co-operation and Development
SMR	standardised mortality ratio

METHODOLOGY FOR MEASURING IN-HOSPITAL MORTALITY

MORTALITY RATES

Introduction

In-hospital mortality (death) rates measure the number of deaths as a proportion of the number of hospital admissions. Differences in mortality findings between hospitals can be due to one or more of the following:

- **Expected variation:** Due to the nature of data, there will always be some fluctuation in the precise measure between one reporting period and the next.
- **Differences in patient factors:** Patients differ from one another in terms of age, gender and comorbidities.
- Differences in data collection: Hospitals and healthcare providers may for many complex reasons differ in terms of how a patient's medical chart is completed or how conditions are recorded and coded.
- **Differences in quality of care:** Mortality is one measure of the quality of care provided to patients in hospital.

There are a number of approaches to measuring mortality rates. Each varies; they are calculated in different ways and used for different purposes. The three main approaches are:

- 1. Crude in-hospital mortality rate
- 2. Directly standardised in-hospital mortality rate
- 3. Indirectly standardised in-hospital mortality ratio.

1. Crude in-hospital mortality rate

The crude in-hospital mortality rate is a measure of the number of deaths per 100 admissions. It is important to remember that it does not attempt to adjust for differences in patient populations (such as age and comorbidity; see above). It is usually presented with reference to a specific disease, such as stroke or acute myocardial infarction (AMI). It is typically expressed as the number of deaths per 100 of the total number of admissions for that specific condition per year.


Crude in-hospital mortality rate for a given diagnosis *For that specific diagnosis

The crude in-hospital mortality rate gives an overview of the extent to which a given condition adds to the overall burden from death in a particular hospital or group of hospitals. It is not a standardised measure because it does not take into consideration confounding factors such as age, type of admission, previous admissions or existing background illness (case mix and comorbidity) in a population of patients. This method is useful in that it allows each hospital to take a bird's-eye view of its number of deaths as well as time trends, providing there has been no significant change in case mix during the period in question.

However, that being said, it is not appropriate to compare hospitals against one another using the crude in-hospital mortality rate because it does not take into account any of the other important factors affecting mortality. Crude in-hospital mortality is used in this report to show the national trend.

2. Directly standardised mortality rate

The direct standardisation method provides more adjustment for population differences. Standardisation in this context means that a common age-structured population is used as the standard, and study populations or groups are compared against this. Age and gender are the two most common variables used for direct standardisation, and the national population may be used as the 'standard population' (see Figure 1).



Standardisation therefore means that the mortality rates produced for a population or condition in a hospital are those that they would have had if they had the same attributes as the standard population. Direct standardisation methods are more powerful when numbers are larger, and are best used for a single or otherwise homogeneous group of diagnoses. It is important to note that only a limited number of variables may be standardised for using this method.

The Organisation for Economic Co-operation and Development (OECD) uses the directly standardised death rate as the basis for its methodological approach (Organisation for Economic Co-operation and Development, 2015). The reference population is based on the age and gender profile of the OECD 2010 population admitted to hospital with selected conditions. This allows direct comparison between OECD member states and is of greatest value when it is used to compare practices across international boundaries. This is the approach used by the Department of Health for the *National Healthcare Quality Reporting System Annual Report* (Department of Health, 2018) for selected diagnoses, specifically AMI, haemorrhagic stroke and ischaemic stroke. Due to the differences in methodology, it is not possible to compare in-hospital mortality indicators in this report against those reported by the Department of Health.

3. Indirectly standardised mortality ratio

The standardised mortality ratio (SMR) is another method that adjusts for population differences. It is a measure of mortality which allows individual hospitals to compare their observed death rate against the rate that would be expected in that hospital if other variables affecting mortality could be taken into consideration. SMRs are analysed based on the principal diagnosis of the patient recorded in HIPE; that is, the diagnosis which was established after investigation and found to be responsible for the episode of admitted patient care. It does not follow that the principal reason for a person's hospitalisation is always the reason for their death.

In equation format:

Standardised mortality ratio =

 $\frac{\text{Observed deaths}}{\text{Expected deaths}} \qquad \text{x 100}$

NATIONAL AUDIT OF HOSPITAL MORTALITY SUPPORTING APPENDIX 2018

The 'expected' deaths are calculated from national data using statistical techniques to account for differences in patient factors. These factors include: age, deprivation, whether patients were in receipt of palliative care treatment in hospital, number of previous admissions in the past year, source and type of admission (for example, from home or a nursing home or an emergency transfer from another acute hospital), and the Charlson Comorbidity Index (Charlson *et al.*, 1987), which is a measure of comorbidity. The Charlson Comorbidity Index assigns a weighting to the degree to which the patient is debilitated by a number of background illnesses and conditions. Therefore, it can be seen that the indirect approach to standardisation allows a greater number of variables to be controlled for, which is very useful in dealing with complex conditions and presentations.

Overall, the SMR is an appropriate way to measure in-hospital mortality in Ireland because:

- There are a large number of hospitals, many of very different sizes.
- It takes account of a larger number of variables, which impact on in-hospital mortality.

SMRs can be presented by individual hospital or by diagnosis group, such as AMI or stroke. They do not allow hospitals to compare outcomes against one another, but they do allow comparison against a national average, which is set at 100.

Control limits

While the national average is set at 100, it is unlikely that any calculated SMR will be exactly 100. A certain amount of variation above or below this average is to be expected as normal, and the control limit approach is a method of accounting for this. Control limits are statistical calculations based on the number of admissions and deaths within each hospital which show the variation that is normally expected to occur in that hospital's data. The control limits are set at 99.8%, meaning that there is a 1-in-500 chance of a hospital being outside these limits by chance alone. This means, therefore, that an SMR which is above (or below) the 99.8% control limits is unlikely to have occurred by chance and may indicate greater (or fewer) deaths than would otherwise be expected.

References

Charlson, M.E., Pompei, P., Ales, K.L. and MacKenzie, C.R. (1987). A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *Journal of Chronic Diseases* 40(5): pp. 373-383. Available from: http://www.aqc.ch/download/HSM_Suppl_8_charlson.pdf [Accessed 12 August 2019]

Department of Health (2018). *National Healthcare Quality Reporting System: Annual Report 2018*. Available from: https://health.gov.ie/wp-content/uploads/2018/07/2018-NHQRS-Report-1.pdf [Accessed 12 August 2019]

Naing, N.N. (2000). Easy Way to Learn Standardization: Direct and Indirect Methods. *Malaysian Journal of Medical Sciences*, 7(1): pp. 10-15. Available from: https://www.ncbi. nlm.nih.gov/pmc/articles/PMC3406211/ [Accessed 12 August 2019]

Organisation for Economic Co-operation and Development (2015). *Definitions for Health Care Quality Indicators 2014-2015 HCQI Data Collection*. Available from: http://stats.oecd. org/wbos/fileview2.aspx?IDFile=62f94ae6-180c-4e4b-9a22-b030ddadfd35[Accessed 12 August 2019]

FRAMEWORK FOR THE NAHM ANNUAL REPORT

TABLE 1: CRITERIA FOR SELECTION OF KEY DIAGNOSES

The National Audit of Hospital Mortality (NAHM) Governance Committee defined inclusion criteria in order to select a cohort of key diagnoses for the National Audit of Hospital Mortality Annual Report, as shown in Table 1.

	Criterion	Comment	Rationale
	Alignment to National Clinical Programme	Is there an aligned Health Service Executive (HSE) National Clinical Programme?	HSE National Clinical Programmes provide national leadership for improvement.
Clinical	Burden of the clinical topic	Is the key diagnosis considered of high volume?	HSE National Clinical Programmes provide national leadership for improvement.
	Significant clinical risk	Is the key diagnosis considered of significant clinical risk, for example high mortality?	
Methodological	Definition	Is the volume of expected deaths ≥5?	Only key diagnoses which are explicitly defined are selected for reporting.
	Number of hospitals with defined number of admissions and expected events	Is the volume of expected deaths ≥5? Is the volume of admissions >100 over the reporting period for the individual diagnosis?	The model is more statistically reliable when these criteria are met.
	Statistical validity of the model	Is the receiver operating characteristic (ROC) statistic >0.7?	This measure calculates the performance of the model in predicting death. A result of >0.7 is considered a satisfactory predictor.

NAHM GOVERNANCE STRUCTURE

NAHM is deployed under the governance framework of the National Office of Clinical Audit (NOCA). The NOCA Governance Board guides NOCA's clinical decision-making and strategic direction, and it provides oversight for seven national clinical audits.

NOCA has established a NAHM Governance Committee with multidisciplinary membership, including clinical and executive leadership from Irish hospitals and the health service. The NAHM Governance Committee oversees two subcommittees and the NOCA Operational Team.

- In consultation with the Health Intelligence, Strategic Planning and Transformation, Health Service Executive (HIU), NOCA has developed the NQAIS NAHM Analysis and Display Scientific Team (ADST). This team provides specialist expertise in order to achieve excellence in the development and enhancement of the National Quality Assurance Improvement System National Audit of Hospital Mortality (NQAIS NAHM) web-based tool. The membership of the team comes from the HIU, NOCA, the NAHM Governance Committee and the software developer. This aligns NQAIS NAHM with developments on other NQAIS projects under the HIU umbrella.
- The NAHM Writing Group is a subcommittee of the NAHM Governance Committee that is convened specifically to write the annual report.
- The NAHM Operational Team oversees the day-to-day management of NAHM in line with NOCA policies and guidelines.




TABLE 2: ATTENDANCE AT NAHM GOVERNANCE COMMITTEE MEETINGS, 2018

		Date	Meeting	Meeting	Meeting	Meeting	Meeting	
Name	Role	Joined	24/01/18	10/04/18	05/07/18	11/09/18	16/10/18	Comment
Margaret Brennan	HSE Acute Operations Division	2016	1	1	Р	Ρ	Ρ	Gareth Clifford Proxy
Ronan Collins	Clinical Programmes Representative	2018	N/A	\bowtie	1	\bowtie		Joined April 2018
Brian Creedon	Chair; Royal College of Physicians of Ireland (Pal Care)	2015	1	1	1	1	1	Brian retired following launch of annual report 2017
Bridget Egan	Royal College of Surgeons in Ireland	2015	1	1	1	1	\bowtie	Appointed as Clinical Lead from January 2019
Alan Egan	Patient and Public Involvement Representative	2017	1	1	1	1	\bowtie	
Eilísh Hardiman	HSE Hospital Groups Forum	2015	1	1	1	1	1	
Ray Healy	NAHM User Representative	2018	N/A	N/A	N/A	1	1	Joined July 2018, Retired December 2018
Howard Johnson	HSE Health Intelligence Unit	2015	1	1	1	1	1	
Simon Jones	International expert	2015	1	1	\bowtie	1	1	
Niall Mahon	Royal College of Physicians of Ireland (Cardiology)	2015	\bowtie	N/A	N/A	N/A	N/A	Retired January 2018
Jennifer Martin	HSE Quality Improvement Team	2016	\bowtie	Ρ	1	1	Ρ	Appointed as Chair from January 2019
Jeanne Moriarty	Joint Faculty of Intensive Care Medicine of Ireland	2018	1	\bowtie	\bowtie	\bowtie	1	
Ed McKone	Royal College of Physicians of Ireland (Resp Med)	2017	\bowtie		1	\bowtie	\bowtie	Joined April 2017
Deirdre Murphy	HSE Healthcare Pricing Office	2015	1	1	\bowtie	1	1	
Brian O'Mahony	Public Representative	2015			\bowtie	N/A	N/A	Retired
Geraldine Shaw	HSE Office of Nursing and Midwifery Services	2015	\bowtie		1	1		
Collette Tully	NOCA	2016	N/A	N/A	N/A	N/A	N/A	Attendance as agenda requires
Barry White	Royal College of Physicians of Ireland	2015						Retired
Deirdre Burke	NOCA	2015	1	1	1	1	1	
Marinia Cronin	NOCA	2015	\square	1	\square	1	1	

✓ Attended Meeting X Apologies received

P Proxy attending N/A not yet appointed or resigned from committee

INDICATORS FOR KEY DIAGNOSIS AND TABULAR INFORMATION TO SUPPORT FUNNEL PLOTS IN THE MAIN REPORT ACUTE MYOCARDIAL INFARCTION

TABLE 3: AMI INDICATOR

Definition	Standardised mortality ratio with a principal diagnosis of AMI
Year covered	2018
ICD-10-AM codes	121, 1210, 1211, 1212, 1213, 1214, 1219, 122, 1220, 1221, 1228, 1229
Methodology	 Numerator Number of actual deaths following admission to hospital with the following ICD-10-AM principal diagnoses: 'Acute myocardial infarction', 'Acute transmural MI of anterior wall', 'Acute transmural MI of inferior wall', 'Acute transmural MI of other sites', 'Acute transmural MI of unspecified site', 'Acute sub-endocardial MI', 'Acute myocardial infarction unspecified', 'Subsequent myocardial infarction', 'Subsequent MI of anterior wall', 'Subsequent MI of inferior wall', 'Subsequent MI of inferior wall', 'Subsequent MI of inferior wall', 'Subsequent MI of other sites', 'Subsequent MI of unspecified site' Denominator Number of expected deaths for AMI. This is calculated using an indirect standardisation and logistic regression modelling of all patients admitted to hospital with a principal diagnosis of AMI.

Hospital Group	Hospital name	No. of admissions for AMI, 2018	SMR -AMI, 2018	99.8% control limits
Dublin	Naas General Hospital	125	104	(-3-232)
Midlands	St James's Hospital	664	122	(42-166)
Hospital Group	Tallaght University Hospital(Adult)	229	129	(5-216)
	Mater Misericordiae University Hospital	557	146	(33-177)
Ireland	Regional Hospital Mullingar	101	119	(-17-255)
East Hospital	Our Lady's Hospital, Navan	115	0	(-15-250)
Group	St Luke's Hospital, Kilkenny	180	147	(1-224)
	St Vincent's University Hospital	299	131	(27-185)
	Wexford General Hospital	235	62	(18-199)
RCSI	Connolly Hospital	141	60	(-3-233)
Hospitals	Beaumont Hospital	227	82	(1-336)
	Cavan General Hospital	189	79	(24-189)
	Our Lady of Lourdes Hospital, Drogheda	195	65	(24-189)
Saolta	Galway University Hospitals	495	144	(40-167)
University Health Care	Letterkenny University Hospital	210	67	(14-204)
Group	Mayo University Hospital	172	169	(11-209)
	Sligo University Hospital	147	42	(19-197)
South /	Cork University Hospital	661	124	(44-163)
South West Hospital	University Hospital Kerry	137	36	(17-201)
Group	South Tipperary General Hospital	106	113	(-15-250)
	University Hospital Waterford	162	126	(0-228)
UL Hospital Group	University Hospital Limerick	462	76	(39-170)

TABLE 4: TABULAR PRESENTATION FOR AMI IN-HOSPITAL MORTALITY, 2018

HEART FAILURE

TABLE 5: HEART FAILURE INDICATOR

Definition	Standardised mortality ratio with a principal diagnosis of heart failure
Year covered	2018
ICD-10-AM codes	150, 1500, 1501, 1509
Methodology	Numerator Number of actual deaths following admission to hospital with the following ICD-10-AM principal diagnoses: 'Heart failure', 'Congesfve heart failure', 'Leg ventricular failure', 'Heart failure unspecified'
	Denominator Number of expected deaths for heart failure. This is calculated using an indirect standardisation and logistic regression modelling of all patients admitted to hospital with a principal diagnosis of heart failure.

Hospital Group	Hospital name	No. of admissions for Heart Failure, 2018	SMR - Heart Failure	99.8% control limits
Dublin	Naas General Hospital	127	106	(11-210)
Midlands	Midland Regional Hospital, Portlaoise	119	114	(18-199)
Hospital	St James's Hospital	255	141	(38-171)
Group	Tallaght University Hospital (Adult)	168	85	(18-197)
	Midland Regional Hospital, Tullamore	143	124	(18-200
	Mater Misericordiae University Hospital	279	108	(39-170)
Ireland	Regional Hospital Mullingar	169	72	(17-201)
East Hospital	Our Lady's Hospital, Navan	162	118	(21-194)
Group	St Luke's Hospital, Kilkenny	215	137	(31-181)
	St Vincent's University Hospital	332	106	(50-156
	Wexford General Hospital	263	114	(28-183
RCSI	Connolly Hospital	156	97	(30-181)
Hospitals	Beaumont Hospital	276	123	(42-165)
	Cavan General Hospital	201	100	(24-189)
	Our Lady of Lourdes Hospital, Drogheda	229	89	(28-183)
Saolta	Portiuncula University Hospital	110	52	(11-209)
University	Galway University Hospitals	440	102	(49-156)
Health Care	Letterkenny University Hospital	218	101	(29-183)
Group	Mayo University Hospital	345	103	(43-163)
	Sligo University Hospital	189	58	(31-180)
	Bantry General Hospital	115	169	(2-222)
South /	Cork University Hospital	382	112	(36-173)
South West	University Hospital Kerry	154	105	(11-209)
Hospital	Mallow University Hospital	130	140	(0-228)
Group	Mercy University Hospital	164	106	(13-204
	South Tipperary General Hospital	136	30	(12-208)
	University Hospital Waterford	198	53	(27-186)
		20.4	00	
UL Hospital	University Hospital Limerick	294	88	(43-164)
Group	Ennis Hospital	157	59	(31-180)
	St John's Hospital, Limerick	119	81	(24-190)
	Nenagh Hospital	125	81	(1-226)

TABLE 6: TABULAR PRESENTATION FOR HEART FAILURE IN-HOSPITAL MORTALITY, 2018

ISCHAEMIC STROKE

TABLE 7: ISCHAEMIC STROKE INDICATOR

Definition	Standardised mortality ratio with a principal diagnosis of ischaemic stroke
Year covered	2018
ICD-10-AM codes	163, 1630, 1631, 1632, 1633, 1634, 1635, 1636, 1638, 1639
Methodology	 Numerator Number of actual deaths following admission to hospital with the following ICD-10-AM principal diagnoses: 'Cerebral infarction', 'Cerebral infarction due to thrombosis of the pre-cerebral artery', 'Cerebral infarction due to embolism of pre-cerebral artery', 'Cerebral infarction due to unspecified occlusion of pre-cerebral artery', 'Cerebral infarction due to thrombosis of the cerebral artery', 'Cerebral infarction due to embolism of the cerebral artery', 'Cerebral infarction due to embolism of the cerebral artery', 'Cerebral infarction due to unspecified occlusion of us to unspecified occlusion of the cerebral artery', 'Cerebral infarction due to unspecified occlusion of the cerebral artery', 'Cerebral infarction due to unspecified occlusion of the cerebral artery', 'Cerebral infarction due to central venous thrombosis non-pyogenic', 'Other cerebral infarction', 'Cerebral infarction unspecified' Denominator Number of expected deaths for ischaemic stroke. This is calculated using an indirect standardisation and logistic regression modelling of all patients admitted to hospital with a principal diagnosis of ischaemic stroke.

Hospital Group	Hospital name	No. of admissions for Ischaemic Stroke, 2018	SMR – Ischaemic Stroke, 2018	99.8% control limits	
Dublin	Naas General Hospital	177	87	(29-182)	
Midlands	St James's Hospital	229	134	(19-197)	
Hospital Group	Tallaght University Hospital (Adult)	227	154	(20-195)	
Ireland East Hospital	Mater Misericordiae University Hospital	297	86	(42-166)	
Group	St Vincent's University Hospital	321	107	(48-158)	
	Wexford General Hospital	110	169	(8-213)	
RCSI Hospitals	Connolly Hospital Beaumont Hospital Cavan General Hospital Our Lady of Lourdes Hospital, Drogheda	136 501 130 181	105 75 98 76	(6-216) (45-162) (20-195) (31-180)	
Saolta University Health Care Group	Galway University Hospitals Letterkenny University Hospital Mayo University Hospital Sligo University Hospital	256 147 144 124	53 139 91 102	(39-169) (18-199) (25-188) (12-208)	
South / South West Hospital Group	Cork University Hospital University Hospital Kerry	433 161	163 55	(42-165)	
UL Hospital Group	University Hospital Limerick	317	78	(44-163)	

TABLE 8: TABULAR PRESENTATION FOR ISCHAEMIC STROKE IN-HOSPITAL MORTALITY, 2018

HAEMORRHAGIC STROKE

TABLE 9: HAEMORRHAGIC STROKE INDICATOR

Definition	Standardised mortality ratio with a principal diagnosis of haemorrhagic stroke
Years covered	2016-2018
ICD-10-AM codes	160, 1600, 1601, 1602, 1603, 1604, 1605, 1606, 1607, 1608, 1609, 161, 1610, 1611, 1612, 1613, 1614, 1615, 1616, 1618, 1619
Methodology	Numerator Number of actual deaths following admission to hospital with the following ICD- 10-AM principal diagnoses: 'Subarachnoid haemorrhage', 'Subarachnoid haemorrhage, carotid siphon and bifurcation', 'Subarachnoid haemorrhage from middle cerebral artery', 'Subarachnoid haemorrhage from anterior communicating artery', 'Subarachnoid haemorrhage from posterior communicating artery', 'Subarachnoid haemorrhage from basilar artery', 'Subarachnoid haemorrhage from vertebral artery', 'Subarachnoid haemorrhage from other intracranial artery', 'Subarachnoid haemorrhage from intracranial artery unspecified', 'Other subarachnoid haemorrhage', 'Subarachnoid haemorrhage unspecified', 'Intracerebral haemorrhage', 'Intracerebral haemorrhage in hemisphere subcortical', 'Intracerebral haemorrhage in hemisphere cortical', 'Intracerebral haemorrhage in cerebellum', 'Intracerebral haemorrhage in brain stem', 'Intracerebral haemorrhage in cerebellum', 'Intracerebral haemorrhage', 'Intracerebral haemorrhage unspecified', 'Other intracerebral haemorrhage', 'Intracerebral haemorrhage unspecified', 'Other intracerebral haemorrhage', 'Intracerebral haemorrhage unspecified', 'Intracerebral haemorrhage', 'Intracerebral haemorrhage unspecified', 'Other intracerebral haemorrhage', 'Intracerebral haemorrhage unspecified', 'Intracerebral haemorrhage', 'Intracerebral haemorrhage unspecified', 'Other intracerebral haemorrhage', 'Intracerebral haemorrhage unspecified'.
	Denominator Number of expected deaths for haemorrhagic stroke. This is calculated using an indirect standardisation and logistic regression modelling of all patients admitted to hospital with a principal diagnosis of haemorrhagic stroke.

TABLE 10: TABULAR PRESENTATION FOR HAEMORRHAGIC STROKE IN-HOSPITAL MORTALITY, 2016-2018				
Hospital Group	Hospital name	No. of admissions for Haemorrhagic Stroke, 2016-2018	SMR – Haemorrhagic Stroke, 2016-2018	99.8% control limits
Dublin Midlands	Naas General Hospital	102	98	(49-157)
Hospital Group	St James's Hospital	155	120	(53-151)
	Tallaght University Hospital (Adult)	137	89	(50-155)
Ireland East Hospital Group	Mater Misericordiae University Hospital St Vincent's University Hospital	248 283	75 101	(64-139) (71-131)
RCSI Hospitals	Connolly Hospital Beaumont Hospital Our Lady of Lourdes Hospital, Drogheda	107 923 112	102 100 98	(49-156) (73-129) (52-153)
Saolta University Health Care Group	Galway University Hospitals	157	99	(60-143)
South / South West Hospital Group	Cork University Hospital University Hospital Waterford	486 104	126 79	(71-130) (53-152)
UL Hospital Group	University Hospital Limerick	211	110	(63-140)

TABLE 10: TABULAR PRESENTATION FOR HAEMORRHAGIC STROKE IN-HOSPITAL MORTALITY, 2016-2018

COPD

TABLE 11: COPD INDICATOR		
Definition	Standardised mortality ratio with a principal diagnosis of COPD	
Year covered	2018	
ICD-10-AM codes	J40, J41, J410, J411, J418, J42, J43, J430, J431, J432, J438, J439, J44, J440, J441, J448, J449, J47	
Methodology	 Numerator Number of actual deaths following admission to hospital with the following ICD-10-AM principal diagnoses: 'Bronchitis not specified as acute or chronic', 'Simple and mucopurulent chronic bronchitis', 'Simple chronic bronchitis', 'Mucopurulent chronic bronchitis', 'Mixed simple and mucopurulent chronic bronchitis', 'Unspecified chronic bronchitis', 'Emphysema', 'MacLeod's syndrome', 'Pan-lobular emphysema', 'Centrilobular emphysema', 'Other emphysema', 'Emphysema unspecified', 'Other COPD', 'COPD with acute lower respiratory infection', 'COPD with acute exacerbation unspecified', 'Other specified COPD', 'COPD unspecified', 'Bronchiectasis' Denominator Number of expected deaths for COPD. This is calculated using an indirect standardisation and logistic regression modelling of all patients admitted to hospital with a principal diagnosis of COPD. 	

Hospital Group	Hospital name	No. of admissions for COPD, 2018	SMR -COPD, 2018	99.8% control limits
Dublin Midlands Hospital Group	Naas General Hospital	411	106	(33-178)
	Midland Regional Hospital, Portlaoise	337	91	(16-201)
	St James's Hospital	789	96	(48-157)
	Tallaght University Hospital (Adult)	769	131	(45-162)
	Midland Regional Hospital, Tullamore	448	140	(25-188)
	St Columcille's Hospital	260	53	(-13-245
	Mater Misericordiae University Hospital	834	104	(38-171)
reland	Regional Hospital Mullingar	441	87	(21-194)
ast .	Our Lady's Hospital, Navan	390	96	(23-191)
Hospital Group	St Luke's Hospital, Kilkenny	548	85	(38-171)
broap	St Michael's Hospital	193	17	(-11-241
	St Vincent's University Hospital	415	84	(32-178
	Wexford General Hospital	763	111	(40-168
	Connolly Hospital	537	111	(36-174
RCSI	Beaumont Hospital	731	131	(43-164
Hospitals	Cavan General Hospital	506	67	(33-177
	Our Lady of Lourdes Hospital, Drogheda	592	101	(36-174
	Portiuncula University Hospital	307	124	(22-193)
Saolta	Galway University Hospitals	624	67	(41-167)
Jniversity Health Care	Letterkenny University Hospital	621	87	(31-179)
Group	Mayo University Hospital	628	106	(45-161)
	Sligo University Hospital	455	96	(36-174
South /	Cork University Hospital	644	161	(36-173
South / South West	University Hospital Kerry	325	110	(24-190
Hospital	Mallow University Hospital	274	40	(-17-254
Group	Mercy University Hospital	582	124	(37-172)
	South Tipperary General Hospital	406	71	(31-180)
	University Hospital Waterford	314	106	(18-200
UL Hospital Group	University Hospital Limerick	739	108	(54-151)
	Ennis Hospital	368	55	(38-170)
	St John's Hospital, Limerick	444	64	(31-179)
	Nenagh Hospital	171	97	(-16-251

TABLE 12: TABULAR PRESENTATION FOR COPD IN-HOSPITAL MORTALITY, 2018

PNEUMONIA

TABLE 13: PNEUMONIA INDICATOR

Definition	Standardised mortality ratio with a principal diagnosis of pneumonia			
Year covered	2018			
ICD-10-AM codes	A202, A212, A221, A310, A420, A430, A481, A78, B012, B052, B250, B583, B59, B671, J12, J120, J121, J122, J123, J128, J129, J13, J14, J15, J150, J151, J152, J153, J154, J155, J156, J157, J158, J159, J16, J160, J168, J17, J170, J171, J172, J173, J178, J18, J180, J181, J182, J188, J189, J85, J850, J851			
Methodology	Numerator Number of actual deaths following admission to hospital with the following ICD- 10-AM principal diagnoses:			
	 'Pneumonic plague', 'Pulmonary tularaemia', 'Pulmonary anthrax', 'Pulmonary mycobacterial infection', 'Pulmonary actinomycosis', 'Pulmonary nocardiosis', 'Legionnaires' disease', 'Q fever', 'Varicella pneumonia', 'Measles complicated by pneumonia', 'Cytomegaloviral pneumonitis', 'Pulmonary toxoplasmosis', 'Pneumocystosis', 'Echinococcus granulosus infection lung', 'Viral pneumonia not elsewhere classified', 'Adenoviral pneumonia', 'Respiratory syncytial virus pneumonia', 'Parainfluenza virus pneumonia', 'Human metapneumovirus pneumonia', 'Other viral pneumonia', 'Viral pneumonia unspecified', 'Pneumonia due to Streptococcus pneumonia', 'Pneumonia due to Haemophilus influenzae', 'Bacterial pneumonia NEC', 'Pneumonia due to Klebsiella pneumoniae', 'Pneumonia due to Streptococcus group B', 'Pneumonia due to other streptococci', 'Pneumonia due to streptococcus group B', 'Pneumonia due to other streptococci', 'Pneumonia due to Escherichia coli', 'Pneumonia due to other (aerobic) gram negative bacteria', 'Bacterial pneumonia in disease classified elsewhere', 'Pneumonia due to other spec infect organisms', 'Pneumonia in diseases class elsewhere', 'Pneumonia in bacteria disease classified elsewhere', 'Pneumonia in viral disease classified elsewhere', 'Pneumonia unspecified', 'Hypostatic pneumonia unspecified', 'Cher pneumonia unspecified', 'Pneumonia unspecified', 'Abscess of lung and mediastinum', 'Gangrene and necrosis of lung', 'Abscess of lung with pneumonia' 			

Number of expected deaths for pneumonia. This is calculated using an indirect standardisation and logistic regression modelling of all patients admitted to hospital with a principal diagnosis of pneumonia.

Hospital Group	Hospital name	No. of admissions for pneumonia, 2018	SMR – pneumonia, 2018	99.8% control limits
Dublin	Naas General Hospital	404	81	(56-148)
Midlands	Midland Regional Hospital, Portlaoise	238	79	(43-165)
Hospital	St James's Hospital	854	137	(68-134)
Group	Tallaght University Hospital (Adult)	561	104	(61-142)
	Midland Regional Hospital, Tullamore	236	168	(37-172)
	St Columcille's Hospital	172	43	(18-197)
	Mater Misericordiae University Hospital	803	97	(64-139
Ireland	Regional Hospital Mullingar	240	108	(42-165)
East	Our Lady's Hospital, Navan	352	61	(53-152)
Hospital Group	St Luke's Hospital, Kilkenny	502	91	(60-143
Group	St Michael's Hospital	214	87	(41-166)
	St Vincent's University Hospital	848	96	(71-131)
	Wexford General Hospital	341	87	(44-163
RCSI Hospitals	Connolly Hospital Beaumont Hospital Cavan General Hospital	484 514 507	113 136 93	(61-142) (58-145) (60-144
	Our Lady of Lourdes Hospital, Drogheda	862	95	(69-133)
Saolta	Portiuncula University Hospital	520	72	(63-140)
University	Galway University Hospitals	616	99	(62-141)
Health Care	Letterkenny University Hospital	430	119	(56-148
Group	Mayo University Hospital	407	127	(54-150
	Sligo University Hospital	646	68	(66-136
	Bantry General Hospital	103	139	(18-198)
South /	Cork University Hospital	518	165	(55-149)
South West	University Hospital Kerry	396	82	(57-147)
Hospital	Mallow University Hospital	176	124	(26-187)
Group	Mercy University Hospital	304	137	(46-161)
	South Tipperary General Hospital	380	95	(57-147)
	University Hospital Waterford	274	87	(48-158
UL Hospital Group	University Hospital Limerick	408	90	(54-150)
	Ennis Hospital	140	53	(34-176)
	St John's Hospital, Limerick	162	50	(38-170)

TABLE 14: TABULAR PRESENTATION FOR PNEUMONIA IN-HOSPITAL MORTALITY, 2018



Phone: +353 1 4028577 Email: nahm@noca.ie Twitter: @noca_irl

www.noca.ie

