

## Effect of essential oils of *Syzygium aromaticum* and *Cinnamomum zeylanicum* and their major components on biofilm production in *Staphylococcus aureus* strains isolated from milk of cows with mastitis.

### AUTHOR(S)

Paulo E. Budri, Nathalia CC Silva, Erika CR Bonsaglia, Ary Fernandes Júnior, Joao P. Araújo Júnior, Julio T. Doyama, Juliano L. Gonçalves, M V. Santos, Deirdre Fitzgerald-Hughes, Vera LM Rall

### CITATION

Budri, Paulo E.; Silva, Nathalia CC; Bonsaglia, Erika CR; Júnior, Ary Fernandes; Júnior, Joao P. Araújo; Doyama, Julio T.; et al. (2015): Effect of essential oils of *Syzygium aromaticum* and *Cinnamomum zeylanicum* and their major components on biofilm production in *Staphylococcus aureus* strains isolated from milk of cows with mastitis.. Royal College of Surgeons in Ireland. Journal contribution.  
<https://hdl.handle.net/10779/rcsi.10768223.v1>

### HANDLE

[10779/rcsi.10768223.v1](https://hdl.handle.net/10779/rcsi.10768223.v1)

### LICENCE

CC BY-NC-SA 4.0

This work is made available under the above open licence by RCSI and has been printed from <https://repository.rcsi.com>. For more information please contact [repository@rcsi.com](mailto:repository@rcsi.com)

### URL

[https://repository.rcsi.com/articles/journal\\_contribution/Effect\\_of\\_essential\\_oils\\_of\\_Syzygium\\_aromaticum\\_and\\_Cinnamomum\\_zeylanicum\\_and\\_their\\_major\\_components\\_on\\_biofilm\\_production\\_in\\_Staphylococcus\\_aureus\\_strains\\_isolated\\_from\\_milk\\_of\\_cows\\_with\\_mastitis\\_/10768223/1](https://repository.rcsi.com/articles/journal_contribution/Effect_of_essential_oils_of_Syzygium_aromaticum_and_Cinnamomum_zeylanicum_and_their_major_components_on_biofilm_production_in_Staphylococcus_aureus_strains_isolated_from_milk_of_cows_with_mastitis_/10768223/1)

**Running title:** Effect of natural products on biofilm production

**Effect of essential oils of *Syzygium aromaticum* and *Cinnamomum zeylanicum* and their major components on biofilm production in *Staphylococcus aureus* strains isolated from milk of cows with mastitis**

Budri, P. E. <sup>\*†</sup>; Silva, N. C. C. <sup>‡</sup>; Bonsaglia, E. C. R. <sup>\*</sup>; Fernandes Júnior, A. <sup>\*</sup>; Araújo Júnior, J.P. <sup>\*</sup>; Doyama, J. T. <sup>§</sup>; Gonçalves, J.L. <sup>#</sup>, Santos, M. V. <sup>#</sup>; Fitzgerald-Hughes D.<sup>†</sup>, Rall, V. L. M. <sup>\*</sup>;

<sup>\*</sup> Department of Microbiology e Immunology, Instituto de Biociências, UNESP – Universidade Estadual Paulista. Botucatu-SP, Brazil.

<sup>†</sup> Department of Clinical Microbiology, Royal College of Surgeons in Ireland, RCSI, Beaumont Hospital, Dublin, Ireland.

<sup>‡</sup> Department of Food Science, Faculty of Food Engineering (FEA), University of Campinas (UNICAMP), Campinas, Sao Paulo, Brazil.

<sup>§</sup>Department of Chemistry e Biochemistry, Instituto de Biociências, UNESP – Universidade Estadual Paulista. Botucatu-SP, Brazil.

<sup>#</sup>Departament of Nutrition and Animal Production, Faculdade de Medicina Veterinária e Zootecnia, USP- Universidade de São Paulo

Corresponding author: Vera Lúcia Mores Rall, Microbiology and Immunology Department, Instituto de Biociências, UNESP – Universidade Estadual Paulista (UNESP), Rubião Junior/Botucatu, CEP 18618-000, Botucatu, São Paulo/Brasil  
vlmores@ibb.unesp.br

## **Interpretative Summary**

The production of biofilm by *S. aureus* is a problem in industry because this could facilitate the adhesion of bacteria to solid surfaces and contributes to the transmission of these bacteria during the food production. This work aims to investigate the effect of the essential oils of *Syzygium aromaticum* (clove) (EOSA) and *Cinnamomum zeylanicum* (cinnamon) (EOCZ) and their major components, eugenol and cinnamaldehyde, on *S. aureus* biofilm formation on different surfaces. The activity of these substances on stainless steel and polystyrene indicates their potential for an alternative sanitizing spray for surface cleaning in the food processing industry.

## Abstract

Bovine mastitis is an inflammation of the mammary glands of cows and causes significant economic losses in dairy cattle. *Staphylococcus aureus* is one of the microorganisms most commonly isolated. Novel agents are required in agricultural industries to prevent the development of mastitis. The production of biofilm by *S. aureus* facilitates the adhesion of bacteria to solid surfaces and contributes to the transmission and maintenance of these bacteria. The effect of the essential oils of *Syzygium aromaticum* (clove) (EOSA) and *Cinnamomum zeylanicum* (cinnamon) (EOCZ) and their major components, eugenol and cinnamaldehyde, on *S. aureus* biofilm formation on different surfaces was investigated. The results showed a significant inhibition of biofilm production by EOSA on polystyrene and stainless steel surfaces (69.4% and 63.6%, respectively). However, its major component, eugenol, was less effective on polystyrene and stainless steel (52.8% and 19.6%,  $p>0.05$ ). Both EOCZ and its major component, cinnamaldehyde, significantly reduced biofilm formation on polystyrene (74.7% and 69.6%, respectively) and on stainless steel surfaces (45.3% and 44.9%, respectively). These findings suggest that EOSA, EOCZ and cinnamaldehyde at 0.106 mg.mL<sup>-1</sup> may be considered for applications such as sanitizers in the food industry.

Keywords: mastitis, antibiofilm *Staphylococcus aureus*, *Syzygium aromaticum* and *Cinnamomum zeylanicum*.

Chemical compounds studied in this article:

Eugenol (PubChem CID: 3314) and Cinnamaldehyde (PubChem CID 637511).

## 62. 1. Introduction

63

64 Bovine mastitis is an inflammation of the mammary glands in dairy cattle,  
65 usually caused by bacteria. It leads to significant economic losses due to reduced  
66 milk production, increased use of drugs and animal morbidity and mortality  
67 (Melchior et al. 2006). *Staphylococcus aureus* is one of the most important  
68 causative agents of clinical, subclinical or chronic mastitis (Vasudevan et al.,  
69 2003).

70 *S. aureus* can produce biofilms, complex polysaccharide- or protein-bound  
71 bacterial structures that facilitate adhesion and multiplication of bacteria on  
72 environmental surfaces and on animal tissues. Bacteria in biofilms are resistant to  
73 phagocytosis, antimicrobial agents and disinfectants due to the low diffusion  
74 through the matrix and altered cellular metabolism (Donlan and Costerton, 2002).  
75 These protective features of *S. aureus* biofilms promote colonization of the  
76 mammary epithelium which precedes the establishment of a persistent infection.  
77 (Penadés and Lasa, 2006).

78 *S. aureus* polysaccharide production is mediated by the *ica* cluster  
79 (intercellular adhesin) which contains *icaA*, *icaB*, *icaC* and *icaD* (McKenney et al.  
80 1998). Co-expression of *icaA* and *icaD* results in phenotypic expression of capsular  
81 polysaccharide (Arciola et al. 2001). Another important gene involved in biofilm  
82 production in *S. aureus* is *bap* which encodes biofilm associated protein (Bap).  
83 Bap promotes primary binding to surfaces and intercellular adhesion (Lasa and  
84 Penadés, 2006), but its prevalence is reported to be relatively low (Seo et al,  
85 2008).

Due to the high prevalence of biofilm production among *S. aureus* in mastitis there has been increased investment in industrial equipment disinfection programs that target biofilms (Gibson et al., 1999). Essential oils have antimicrobial and anti-biofilm activity against bacteria, parasites (Alexopoulos et al., 2011), fungi (Mari et al., 2003) and viruses (Bishop, 1995). More recently, the EOs of aromatic spices and medicinal plants have been tested for their activity against biofilms (Kwiecinski et. al. 2009). The EOs target different cellular mechanisms, such as inhibition of peptidoglycan synthesis (Ogunlana et al., 1987), modification of bacterial membrane hydrophobicity (Cox et al., 2000) and modulation of quorum sensing (Gao et al., 2003). Sanitizers based on natural products such as EO with specific *S. aureus* anti-biofilm activity may have applications in the beef-processing industry for enhanced surface or carcass cleaning.

The aim of this study was to evaluate the effect of the essential oil of clove (*Syzygium aromaticum*) (EOSA), cinnamon (*Cinnamomum zeylanicum*) (EOCZ) and their major compounds, eugenol and cinnamaldehyde on biofilm formation on stainless steel and polystyrene, by isolates of *S. aureus* recovered from the milk of cows with subclinical mastitis.

## **2. Material and methods**

### **2.1 Samples and bacterial isolation**

A collection of 64 isolates of *S. aureus*, previously recovered from the milk of cows with subclinical mastitis was tested. Presumptive identification was by Gram stain and further confirmation was made based on catalase, coagulase and DNase production, as described by Koneman et al. (2008). Molecular confirmation was by polymerase chain reaction (PCR) amplification of the species-specific staphylococcal nuclease gene (*nuc*) using primers and PCR conditions as outlined in Table 1.

### **2.2 PCR to detect genes linked to biofilm production**

DNA extraction was performed using the Minispin kit (GE Healthcare) according to the manufacturer's instructions. PCR reactions were performed with each primer pair in a final volume of 25 µl containing 2.5 µl of 10X PCR buffer, 2.0 mM magnesium chloride, 200 mM dNTPs, 1U Taq DNA polymerase (Fermentas), 10 pmol of each primer) and 3 µl of the DNA template. The primers used for detection of *icaA*, *icaD* and *bap* and their properties are listed in Table 1. The PCR conditions were those described in the references provided (Table 1) with reactions performed in a Gene Amp PCR System 9700 (Applied Biosystem). PCR products were detected using 1.5% agarose gel in Tris - boric acid - EDTA (TBE) buffer and developed with Sybr Green (Invitrogen ®). Positive and negative controls for *icaA* and *icaD* were *S. aureus* ATCC 35983 and *S. epidermidis* ATCC 12.228. For *bap*, a positive and sequenced strain was used.

### **2.3. Extraction of EOSA and EOCZ and chemical analysis by gas chromatography-mass spectrometry (GC-MS)**

The EOSA and EOCZ were extracted from *Syzygium aromaticum* and *Cinnamomum zeylanicum* by drag steam distiller (model MA480 - Marconi). Densities were calculated according to Fonseca and Librand (2008). The major compounds of the EOSA and EOCZ, eugenol and cinnamaldehyde were sourced commercially (Sigma-Aldrich, UK<sup>®</sup>).

Chemical characterization was determined by gas chromatography coupled to mass spectrometer (GC-MS) (model QP5050A - Shimadzu) with the use of a CBP-5 capillary column with a 0.25 mm of internal diameter and 0.25  $\mu$ m film thickness. The chromatographic conditions were set according to the essential oil analysed. EOSA and EOCZ were identified by matching their mass spectra to reference compounds in the National Institute of Standards and Technology (NIST), mass spectra library (Maryland, USA).

142

### **2.4. Determination of the MIC**

Susceptibility tests were performed in triplicate for EOSA, EOCZ and their major components using the broth microdilution method and CLSI guidelines (CLSI, 2009). Briefly, inocula of *S. aureus* were prepared to the density of 0.5 McFarland using a densitometer (Densichek, BioMérieux) in 0.85% saline solution. The inocula were further diluted to an approximate concentration of  $10^5$  CFU/ml and incubated with test compounds at concentrations of 0.025%, 0.04%, 0.06%, 0.08%, 0.10%, 0.20%, 0.40%, 0.80%, in final volumes of 200  $\mu$ l of Brain Heart Infusion broth (BHI, Oxoid) supplemented with 0.5% Tween 80. Positive growth controls and sterility controls were included.



Plates were incubated at 35°C for 24h, after which time, 50µl of 0.01% resazurin was added to each well. The MIC was recorded as the lowest concentration of EO/EO components at which no growth was observed, as indicated by a change of colour from blue to pink.

## **2.5 Production of biofilm by *S. aureus* in the presence and absence of the EOs and their major compounds**

Isolates were cultured in tryptone soy broth (TSB, Oxoid), at 37°C for 24h, and diluted to approximately 10<sup>8</sup> CFU/ml. In 200 µl of this dilution was added EO's, eugenol and cinnamaldehyde separately to a final concentration of 0.106 mg.mL<sup>-1</sup> (sub-inhibitory concentration based on, MIC determination). The experiment was carried out in triplicate using 96 well microtiter plate. Control assays were prepared similarly but EO and major components were replaced with sterile TSB. Plates were incubated at 35°C for 48h in a static incubator, washed three times with phosphate buffered saline (PBS) pH 7.4, dried at room temperature (RT) and stained with 1% gentian violet. After a further wash with distilled water, absorbance at 570nm was measured using an ELISA plate reader (Babsystems, Multiskan EX). *S. aureus* ATCC 35983 was used as positive control (biofilm producer) and *S. epidermidis* ATCC 12228, as a negative control for biofilm production (Vasudevan et al., 2003).

### **2.5.2 Effect of EOs and major components on biofilm formation on stainless steel**

Sterile stainless steel coupons with a diameter of 1 cm were deposited at the bottom of a 24 well plate and *S. aureus* overnight cultures were diluted to 10<sup>8</sup>

CFU/ml and 300µL aliquots were added to triplicate wells containing the coupons and incubated at 35°C for 48h, in the absence and presence of each EO and their major components (0.106 mg.mL<sup>-1</sup>). The coupons were transferred to a new plate, washed three times with PBS, pH 7.4 and stained with 1% crystal violet for 15 min and washed a further three times. The biofilm was resuspended in 300µL of glacial acetic acid for 15 min and 200µl was transferred to a microplate and the absorbance measured at 570 nm.

## 2.6. Statistical Analysis

To compare the effect on biofilm production in the absence and presence of EOs and major components, analysis of variance was conducted in a randomized design followed by Tukey test means. *P* value <0.05 was considered significant.

## 3. Results

### 3.1. Genotypic analysis for the presence of *icaA*, *icaD* and *bap*

From 64 isolates of *S. aureus*, 26 (40.6%) were positive for all three biofilm genes investigated (*icaA*, *icaD* and *bap*). The *icaA* gene alone was detected in 85.9% and *icaD* in 84.3% of isolates. Representative amplicons from *bap*, *icaA* *icaD* PCRs were partially sequenced and confirmed in GenBank (GenBank accession numbers AY220730.1, CP006838.1 and JN226155.1 respectively).

### **3.2 Chemical analysis by gas chromatography-mass spectrometry (GC-MS)**

Details of the physicochemical characteristics of EOSA, EOCZ, eugenol and cinnamaldehyde are shown in Table 2. The major component of EOSA was eugenol and of EOCZ, cinnamaldehyde (86.59%). MIC values towards *S. aureus* isolates are also shown.

### **3.3 Minimum inhibitory concentration of EOs and major compounds**

The minimum inhibitory concentration of eugenol and EOSA were 0.392 mg.mL<sup>-1</sup> and 0.237 mg.mL<sup>-1</sup>, respectively. The EOCZ and cinnamaldehyde showed MICs of 0.243 mg.mL<sup>-1</sup> and 0.199 mg.mL<sup>-1</sup>.

### **3.4 Effect of EO's and major components on biofilm formation on polystyrene and stainless steel coupons.**

The effect of EOs and their major components on *S. aureus* biofilm formation on polystyrene and stainless steel surfaces are summarised in Table 3. OD values (at 570 nm) obtained in the presence of natural compounds are shown in addition to the percentage remaining following exposure to the natural compounds. The values obtained in the absence of EO or their major components were 0.415 (100%) for polystyrene and 1.07 (100%) in stainless steel.

There were significant differences in biofilm production between isolates grown in the absence and presence of EOs/major components. There was a statistically significant reduction in biofilm formation in the presence of EOSA and EOCZ ( $p < 0.01$ ) on both polystyrene as stainless steel. Cinnamaldehyde and eugenol resulted in a statistically significant reduction in biofilm formation on polystyrene ( $p < 0.001$ ) but on stainless steel, a significant reduction in biofilm was seen for cinnamaldehyde ( $p < 0.01$ ) but not eugenol.

Comparing the anti-biofilm activity of each EO to its major component, similar anti-biofilm activity was found for EOCZ and its major, cinnamaldehyde, on both test surfaces. However, EOSA was more effective in reducing biofilm on polystyrene and stainless steel, than was eugenol.

#### 4. Discussion

Among the 64 isolates, 40.6% carried *icaA*, *icaD* and *bap*, simultaneously. *icaD* and *icaA* are reported more frequently in *S. aureus* isolates from cows with mastitis than *bap* (Atshan and Shamsudin, 2011). However, the prevalence of *bap* among *S. aureus* isolates from bovine mastitis found here are significantly higher than previously reported by others (Vautor et al 2008, Seixas et al., 2014). Cucarella et al. (2001) found 5% positivity for *bap* in 350 strains tested and Vautor et al. (2008) could not detect *bap* among 262 *S. aureus* associated with different diseases and recovered from humans and animals. The relatively high *bap* carriage rate found here compared to other studies, suggests that *bap* acquisition by *S. aureus* may be a recent event or that its transmission by horizontal transfer remains limited despite its presence on a mobile transposon-like element, SaPIbov2. Transmission events involving this gene may be on the increase.

The chemical composition of EO's extracted from *Syzygium aromaticum* and *Cinnamomum zeylanicum* were similar to that found by other authors. The main components of the EOSA were eugenol (90.21%) and eugenol acetate (6.5%). Bauer (2001) observed the same components at concentrations of 75-85% and 8-15% respectively. The components of EOCZ were cinnamaldehyde (86.5%) and benzaldehyde (4.2%) similar to those reported by Unlu et al. (2010), 68.9% and

9.9% respectively. However, unlike the study of Unlu et al, small quantities of cinnamaldehyde acetate (7.4%), were found in the present study. As suggested by Burt et al. (2004), it is possible that intraspecies variations in composition may be due to genetic variation, seasonality, geographic location, harvest time and plant parts used in the preparation of oil.

The MIC values of OE's and their major components towards planktonic cells of *S. aureus* were similar to those observed in previous studies (Unlu, 2010.). From the values of MIC, sub-MIC dose of 0.106 mg.mL<sup>-1</sup> of EO and their component were used for biofilm studies. Furthermore, no corrosion of polystyrene plate surface, attributed to effects of OEs (which may have resulted in false-positives due to dye uptake) was observed at this concentration.

Biofilm formation on polystyrene surfaces decreased by 69.4% and 74% in the presence of EOSA and EOCZ. Cinnamaldehyde also reduced biofilm on polystyrene by 69.6% and eugenol treatment resulted in a 52.8% reduction. The greatest reduction in biofilm formation on stainless steel was found for EOSA (63.5%) followed by EOSZ (45.3) and cinnamaldehyde (44.9%). Eugenol had the least effect on biofilm on this surface (19.6%).

Our finding that the eugenol component of EOSA did not significantly contribute to the anti-biofilm activity of EOSA, suggests that other component within the oil may be responsible. According to Bassolé et al. (2010), the antibacterial activity of an essential oil, is mainly due to its major components, but antimicrobial synergy may result from interactions with other minor components. On the other hand, based on the similarity in biofilm disruption on two surface types, for EOCZ and its major component cinnamaldehyde, we suggest the efficiency of EOCZ is closely linked to its major component.

Several studies have investigated antibacterial activity of natural compounds and several mechanisms have been suggested including, membrane disruption, increased bacterial permeability and leakage of cellular contents and coagulation of cytoplasmic components (Lambert et al., 2001; Ultee and Smid, 2001). Inactivation of membrane proteins specifically may contribute to disruption of biofilm production in the early stages (Ultee et al., 1999). These effects would likely result in loss of adhesion and adsorption to the surface leading to a reduction in biofilm.

Other essential oils and their major components also exhibit inhibition of initial biofilm formation. Carvacrol, one of the main antibacterial components of oregano oil and other essential oils, was reported to inhibit biofilms of *S. aureus* and *S. typhimurium* in the initial growth phase and prevented the formation of mature biofilms (Knowles, et al., 2005). Nostro et al. (2007) reported low biofilm formation by various strains of *S. aureus* and *S. epidermidis* in the presence of sublethal concentrations of EO of oregano, carvacrol and thymol. Similar results were obtained for strains of *S. typhimurium* in the presence of sub-lethal concentrations of EO of thyme, oregano and carvacrol (Soni et al., 2013).

Strong anti-biofilm was observed here for EOSA, EOCZ and cinnamaldehyde against *S. aureus* recovered from cases of sub-clinical bovine mastitis. However, some studies have shown that low concentrations of EOs, like tea tree oil and cinnamaldehyde, can increase bacterial metabolic activity in relation to biofilm production due to environmental stress in *S. aureus* and *Pseudomonas aeruginosa*, respectively (Kwiecinski, 2009).

305. **5. Conclusion**

301        Despite the low extraction yield of EOSA, EOCZ and cinnamaldehyde from  
302        *Syzygium aromaticum* and *Cinnamomum zeylanicum*, the strong anti-biofilm  
303        activity reported here on stainless steel and polystyrene, at low concentrations of  
304        these substances, indicates their potential for development. A specific application  
305        could be as an alternative sanitizing spray for surface cleaning in the food  
306        processing industry.

307

308. **6. Acknowledgements**

309        To Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and  
310        the Conselho Nacional de Pesquisa (CNPq), for financial support.

311

312

313

314

## 3157. 7. References

316

317 Alexopoulos, A., A.C. Kimbaris, S. Plessas, I. Mantzourani, I. Theodoridou, E.  
318 Stavropoulou, M.G. Polissiou, E. Bezirtzoglou. 2011. Antibacterial activities of  
319 essential oils from eight Greek aromatic plants against clinical isolates of  
320 *Staphylococcus aureus*. Anaerobe, 17:1-4.

321 Arciola, C.R., L. Baldassarri, L. Montanaro. 2001. Presence of *icaA* and *icaD*  
322 genes and slime production in a collection of staphylococcal strains from catheter-  
323 associated infections. J.Clin. Microbiol. 39:2151–2156.

324 Atshan, S.S.; M.N. Shamsudin. 2011. Evaluation of phenotypic and genotypic  
325 detection methods for biofilm-forming methicillinsensitive and methicillin-resistant  
326 *Staphylococcus aureus* clinical isolates. Ann. Microbiol. 61:825-831.

327 Bassolé, I.H.M., A. Lamien-Meda, B. Bayala, S. Tirogo, C. Franz, J. Novak, R.C.  
328 Nebié, M.H. Dicko. 2010. Composition and Antimicrobial Activities of *Lippia*  
329 *multiflora* Moldenke, *Mentha x piperita* L. and *Ocimum basilicum* L. Essential Oils  
330 and Their Major Monoterpene Alcohols Alone and in Combination. Molecules  
331 15:7825-7839;

332 Bauer, K., D. Garbe, H. Surburg. 2001. Common Fragrance and Flavor Materials:  
333 Preparation, Properties and Uses. 4<sup>a</sup> edição. Wiley-VCH, Weinheim p. 293.

334 Bishop, C.D. 1995. Antiviral activity of the essential oil of *Melaleuca alternifolia*  
335 (Maiden and Betche) Cheel (tea tree) against tobacco mosaic virus. J. Essent. Oil  
336 Res. 7:641 – 644.

337 Burt, S. 2004. Essential oils: their antibacterial properties and potential applications  
338 in foods—a review. Int. J. Food Microbiol. 94:223- 253.

339 CLSI. 2009. Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria  
340 That Grow Aerobically; Approved Standard - Eighth Edition. *CLSI M7-A8*,



341 Cox, S.D., C.M. Mann, J.L. Markham, H.C. Bell, J.E. Gustafson, J.R. Warmington,  
 342 S.G. Wyllie. 2000. The mode of antimicrobial action of the essential oil of  
 343 *Melaleuca alternifolia* (tea tree oil). J. Appl. Microbiol. 88:170–175.

344 Cucarella, C., C. Colano, J. Valle, B. Amorena, I. Lasa, P. Penades. 2001. Bap, a  
 345 *Staphylococcus aureus* surface protein involved in biofilm formation. J. Bacteriol.  
 346 183:2888–2896.

347 Donlan, R.M., J.W. Costerton. 2002. Biofilms: Survival Mechanisms of Clinically  
 348 Relevant Microorganisms. Clin. Microbiol. Rev. 15:167-193.

349 Fonseca, P., A.P.L. Librand. 2008. Evaluation of physico-chemical and  
 350 phytochemical characteristics of differentt tientes of barbatimão  
 351 (*Stryphnodendron barbatimam*). Braz. J. Pharm. Sci. 44:271-77.

352 Gao, M., M. Teplitski, J. B. Robinson, W. D. Bauer. 2003. Production of substances  
 353 by *Medicago truncatula* that affect bacterial quorum sensing. Mol. Plant-Microbe  
 354 Interact. 16:827–834.

355 Gibson, H.; J.H. Taylor, K. E. Hall, J. T. Holah. 1999. Effectiveness of cleaning  
 356 techniques used in the food industry in terms of the removal of bacterial biofilms. J.  
 357 Appl. Microbiol. 87:41–49.

358 Karpouhtsis, I.; E. Pardali, E. Feggou, S. Kokkini, Z.G. Scouras, P. Mavragani-  
 359 Tshipidou. 1998. Insecticidal and genotoxic activities of oregano essential oils. J.  
 360 Agricult. Food Chem. 46:1111 – 1115.

361 Knowles, J.R., S. Roller, D.B. Murray, A.S. Naidu. 2005. Antimicrobial action of  
 362 carvacrol at different stages of dual-species biofilm development by  
 363 *Staphylococcus aureus* and *Salmonella enterica* serovar Typhimurium. Appl.  
 364 Environm. Microbiol. 71:797-803.

365 Koneman, E. W., S. D. Allen, W. M. Janda, P. C. Schreckenberger, W. C. Winn Jr.  
 366 2008. *Diagnostico Microbiológico—Texto e Atlas Colorido*. 6th ed. Editora  
 367 Guanabara Koogan, Rio de Janeiro, Brazil.

368 Kwiecinski, J., S. Eickb, K.Wojcika. 2009. Effects of tea tree (*Melaleuca alternifolia*)  
 369 oil on *Staphylococcus aureus* in biofilms and stationary growth phase. *Int. J.*  
 370 *Antimicrob. Agents* 33:343–347.

371 Lambert, R.J.W., P.N. Skandamis, P. Coote, G. J.E. Nychas. 2001. A study of the  
 372 minimum inhibitory concentration and mode of action of oregano essential oil,  
 373 thymol and carvacrol. *J. Appl. Microbiol.* 91:453–462.

374 Lasa, I., J.R. Penades. 2006. Bap: a family of surface proteins involved in biofilm  
 375 formation. *Res. Microbiol.* 157:99–107.

376 Mari, M., P. Bertolini, G.C. Pratella. 2003. Non-conventional methods for the  
 377 control of post-harvest pear diseases. *J. Appl. Microbiol.* 94:761–766.

378 Melchior, M.B., H. Vaarkamp, J. Fink-Gremmels. 2006. Biofilms: a role in recurrent  
 379 mastitis infections? *The Vet. J.* 171:398–407.

380 Mckenney, D., J. Hübner, E. Muller, WANG, Y.; GOLDMANN, D.A.; PIER, G.B.  
 381 1998. The *ica* locus of *Staphylococcus epidermidis* encodes production of the  
 382 capsular polysaccharide/adhesin. *Infect. Immun.* 66(10):4711-4720.

383 Nostro, A., R.A. Sudano, G. Bisignano, A. Marino, M.A. Cannatelli, F.C. Pizzimenti.  
 384 2007. Effects of oregano, carvacrol and thymol on *Staphylococcus aureus* and  
 385 *Staphylococcus epidermidis* biofilms. *J. Med. Microbiol.* 56:519-523.

386 Ogunlana, E.O., S. Hoeglund, G. Onawunmi, O. Skoeld. 1987. Effects of  
 387 lemongrass oil on the morphological characteristics and peptidoglycan synthesis of  
 388 *Escherichia coli* cells. *Microbios* 50:43–59.

389 Schalm, O.W., D.D. Noorlander. 1957. Experiments and observations leading to  
 390 development of the California Mastitis Test. J. Am. Vet. Med. Assoc. 130:199-204.

391 Seixas, R., J.P. Santos, R. Bexiga, L.C. Vilela, M. Oliveira. 2014. Antimicrobial  
 392 resistance and virulence characterization of methicillin-resistant staphylococci  
 393 isolates from bovine mastitis cases in Portugal. J. Dairy Sci. 97 (1):340-344.

394 Seo,Y.S., D.Y.Lee, N. Rayamahji, M.L. Kang, H.S. Yoo. 2008. Biofilm-forming  
 395 associated genotypic and phenotypic characteristics of *Staphylococcus* spp.  
 396 isolated from animals and air. Res.Vet. Sci. 85:433–438

397 Soni, K.A.; A. Oladunjoye, R. Nannapaneni, M.W. Schilling, J.L. Silva, B. Mikel.  
 398 2013. Inhibition and inactivation of *Salmonella* Typhimurium biofilms from  
 399 polystyrene and stainless steel surfaces by essential oils and phenolic constituent  
 400 carvacrol. J.Food Protect. v.76(2), p.205-212.

401 Ultee, A., E.P.W. Kets, E.J. Smid. 1999. Mechanisms of action of carvacrol on the  
 402 food-borne pathogen *Bacillus cereus*. Appl. Environm. Microbiol. 65 (10):4606–  
 403 4610.

404 Ultee, A., E.J. Smid. 2001. Influence of carvacrol on growth and toxin production by  
 405 *Bacillus cereus*. Int. J. Food Microbiol. 64:373–378.

406 Unlu, M., E. Ergene, G.V. Unlu, H.S. Zeytinoglu, N. Vural. 2010. Composition,  
 407 antimicrobial activity and *in vitro* cytotoxicity of essential oil from *Cinnamomum*  
 408 *zeylanicum* Blume (Lauraceae). Food Chem. Toxicol. 48:3274–3280.

409 Vancraeynest, D., K. Hermans, F. Haesebrouck. 2004. Genotypic and phenotypic  
 410 screening of high and low virulence *Staphylococcus aureus* isolates from rabbits  
 411 for biofilm formation and MSCRAMMs. Vet. Microbiol. 103:241–247.

412 Vasudevan, P., M.K.M. Nair, T. Annamalai, K.S. Venkitanarayanan. 2003.  
413 Phenotypic and genotypic characterization of bovine mastitis isolates of  
414 *Staphylococcus aureus* for biofilm formation. Vet.Microbiol. 92:179–185.  
415 Vautor, E., G. Abadie, A. Pont, R. Thiery. 2008. Evaluation of the presence of the  
416 bap gene in *Staphylococcus aureus* isolates recovered from human and animals  
417 species. Vet. Microbiol. 127:407–411.  
418  
419

Table 1: Oligonucleotides used in the detection of biofilm genes and confirmation of identification of *Staphylococcus aureus* strains isolated from milk of cows with subclinical mastitis.

gene	primer	sequence	Amplicon size	T <sub>m</sub> (°C)	Reference
<i>bap</i>	bap-F	ccctatatcgaagggtgtagaattg	971 bp	65	Cucarella et al. (2001)
	bap-R	gctgttgaagttaatactgtacctgc			
<i>icaA</i>	<i>icaA</i> -F	cctaactaacgaaggtag	1315 bp	49.5	Vadesuvan et al. (2003)
	<i>icaA</i> -R	aagatatagcgataagtgc			
<i>icaD</i>	<i>icaD</i> -F	aaacgtaagagaggtgg	381 bp	50	Vadesuvan et al. (2003)
	<i>icaD</i> -R	ggcaatatgatcaagatac			
<i>nuc</i>	nuc-F	cgtaaatgcacttgcttcagg	257 bp	55	CRL (UE)
	nuc-R	tcagcaaatgcatcacaacag			

T (°C): annealing temperature, bp: base pairs

425

426 Table 2. Physicochemical analyses and MIC values towards *S. aureus* isolates of EOSA,  
 427 EOCZ and their major compounds.

Species or compound	Density (mg.mL <sup>-1</sup> )	MIC <sup>a</sup> (mg.mL <sup>-1</sup> )	Composition (%)
<i>Syzygium</i> <i>aromaticum</i>	1060	0.392	eugenol (90.2%), eugenol acetate (6.5%), β- caryophyllene (1.3%), others (1.9%)
eugenol	1050	0.237	eugenol (100%)
<i>Cinnamomum</i> <i>zeylanicum</i>	1075	0.243	cinnamaldehyde (86.5%), benzaldehyde (4.2%), cineole (1.7%), cinnamic acid (1.5%) (0.6%), eugenol (0.1%), Others (5.42%)
cinnamaldehyde	1080	0.199	cinnamaldehyde (100%)

428 <sup>a</sup> MIC values shown are the mean found for 26 *S. aureus* isolates; EOSA: clove essential oil; MIC:  
 429 minimum inhibitory concentration; EOCZ: cinnamon essential oil.

430

431

Table 3. *S. aureus* biofilm formation on polystyrene and stainless steel in the presence of EOSA, EOCZ and its major components.

	Polystyrene	stainless steel	p value (pol./ss.)
	OD <sub>570</sub>	OD <sub>570</sub>	
	(% reduction)	(% reduction)	
Control	0.415 ±0,075 (0%) a	1.07 ±0,222 (0%) y	-
EOSA	0.127 ±0,047 (69.4%) b	0.390 ±0,080 (63.5%) z	p<0.01 e p<0.01
Eugenol	0.196 ±0,067 (52.8%) c	0.860 ±0,220 (19.2%) y	p<0,01 e p>0.05
EOCZ	0.105 ±0,040 (74.7%) b	0.585 ±0,158 (45.3%) z	p<0.01 e p<0.01
Cinnamaldehyde	0.126 ±0,078 (69.4%) b	0.589 ±0,100 (44.9%) z	p<0.01 e p<0.01

Values followed by same letter in the column do not differ; EOSA: clove essential oil; EOCZ: cinnamon essential oil; pol: polystyrene; ss: stainless steel